

PRODUCT INFORMATION

Clone ID	Warning: Undefined variable \$hasAttributeValueDescription in C:\wwwroot\mirror\dimabio.com\wp-content\plugins\woocommerce-print-products\publicclass-woocommerce-print-products-public.php on line 2806 DM174
Target	ROR2
Synonyms	ROR2;NTRKR2
Host Species	Rabbit
Description	Anti-ROR2 antibody(DM174); Rabbit mAb
Delivery	In Stock
Uniprot ID	Q01974
IgG type	Rabbit IgG
Clonality	Monoclonal
Reactivity	Human
Applications	ELISA; Flow Cyt; WB
Recommended Dilutions	ELISA 1:5000-10000; Flow Cyt 1:100; WB 1:1000
Purification	Purified from cell culture supernatant by affinity chromatography
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Yefel Storage	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	The protein encoded by this gene is a receptor protein tyrosine kinase and type I transmembrane protein that belongs to the ROR subfamily of cell surface receptors. The protein may be involved in the early formation of the chondrocytes and may be required for cartilage and growth plate development. Mutations in this gene can cause brachydactyly type B; a skeletal disorder characterized by hypoplasia/aplasia of distal phalanges and nails. In addition, mutations in this gene can cause the autosomal recessive form of Robinow syndrome; which is characterized by skeletal dysplasia with generalized limb bone shortening; segmental defects of the spine; brachydactyly; and a dysmorphic facial appearance.
Usage	Research use only
Conjugate	Unconjugated
DIMA Disclaimer	All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement.




Figure 1. ROR2 protein is highly expressed on the surface of HEK293 cell membrane. Flow cytometry analysis with Anti-ROR2 (DM174) on HEK293 cells transfected with human ROR2 (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram), and Isotype antibody on HEK293 transfected with irrelevant protein (Orange histogram).






Figure 2. Flow cytometry analysis of antigen binding of rabbit anti-human ROR2 mAb(DME100174).

(A) DME100174 does not bind to Jurkat cells that do not express ROR2.

(B) A clear peak shift of DME100174 was seen compared to the control when incubated with ROR2-expressing Raji cells, indicating strong binding of DME100174 to ROR2. Antibodies were incubated at 5 µg/mL.



Figure 3. Anti-ROR2 antibody (SKU# DME100174) at 1/1000 dilution

Lane : RAJI(human Burkitt's lymphoma B lymphocyte), whole cell lysate

Secondary : Goat Anti-Rabbit IgG H&L (HRP) at 1/5000 dilution

Predicted band size: 105 kDa

Observed band size: 140 kDa

