

PRODUCT INFORMATION

Warning: Undefined variable ShasAttributeValueDescription in C:\www.root\mirror.dimablo.com\wp-content\plugins\woocommerce-print-products\public.class-woocommerce-print-products-public.php on line 2806 bMC286 Clone ID

SI AMES

CD84;SLAMF5;LY9B;SLAMF5 Synonyme

Host Species Rabbit

Anti-SLAMF5 antibody(DMC286); IgG1 Chimeric mAb Description

Delivery In Stock Uniprot ID Q9UIB8

Rabbit/Human Fc chimeric IgG1 lgG type

Clonality Monoclonal Reactivity Human Flow Cyt Application Recommend Dilutions Flow Cyt 1:100

Purification Purified from cell culture supernatant by affinity chromatography

Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. Formulation & Reconstitution

Storage & Shipping

Scre at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated reczing and thawing). Lyophilized proteins a shipped at ambient temperature and the store of the store

Usage Research use only

Unconjugated
All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverengineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement. DIMA Disclaimer




Figure 1. Flow cytometry analysis with Anti-SLAMF5 (DMC286) on HEK293 cells transfected with human SLAMF5 (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

Email: info@dimabio.com Website: www.dimabio.com

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)






Figure 2. Flow cytometry analysis of antigen binding of anti-human SLAMF5 mAb(DMC100286).
(A) DMC100286 does not bind to CHO-S cells that do not express SLAMF5.
(B) A clear peak shift of DMC100286 was seen compared to the control when incubated with SLAMF5-expressing Raji cells, indicating strong binding of DMC100286 to SLAMF5. Antibodies were incubated at 5 μg/mL.





