

PRODUCT INFORMATION

Warning: Undefined variable ShasAttributeValueDescription in C:\unwwroot\mirror.dimabio.com\wp-content\plugins\woocommerce-print-products\public\class-woocommerce-print-products-public.php on line 2806 fcl Clone ID

TR;TfR;TfR1;Trfr;T9;p90;CD71 Synonyme

Host Species Rabbit

Anti-TFRC antibody(7C1), IgG1 Chimeric mAb Description

Delivery In Stock Uniprot ID P02786

Rabbit/Human Fc chimeric IgG1 lgG type

Clonality Monoclonal Reactivity Human Application Flow Cyt Recommend Dilutions

Purification Purified from cell culture supernatant by affinity chromatography

Formulation & Reconstitution Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Flow Cyt 1/100

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. This gene encodes a cell surface receptor necessay for cellular from uptake by the process of receptor-mediated endocytosis. This receptor is required for erytimopotesis and neurologic development. Multiple alternatively spliced variants have been identified. [provided by RefSeq. Sep 2015] Research use only with the process of receptor-mediated endocytosis. Storage & Shipping Background

Conjugate Unconjugated

All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement. DIMA Disclaimer




Figure 1. Flow cytometry analysis with 1µg/mL Anti-TFRC (7C1) mAb on HEK293 cells transfected with human TFRC (Blue histogram) or HÉK293 transfected with irrelevant protein (Red histogram).

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

Email: info@dimabio.com Website: www.dimabio.com








Figure 2. Flow cytometry analysis of antigen binding of anti-human TFRC mAb(DMC101095). (A) DMC101095 does not bind to CHO-S cells that do not express TFRC. (B) A clear peak shift of DMC101095 was seen compared to the control when incubated with TFRC-expressing Siha cells, indicating strong binding of DMC101095 to TFRC. Antibodies were incubated at 5 μ g/mL.

