

PRODUCT INFORMATION

Common Name	Warning: Undefined variable \$hasAttributeValueDescription in C:\wwwroot\mirror\dimabio.com\wp-content\plugins\woocommerce-print-products\publicclass-woocommerce-print-products-public.php on line 2806 MK-7684
Conjugate	Unconjugated
Synonyms	TIGIT;SIG9;STM3
Applications	ELISA; Flow Cyt
Recommended Dilutions	ELISA 1:5000-10000; Flow Cyt 1:100
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Host Species	Humanized
IgG type	Human IgG1 - kappa
Reactivity	Human
Target	TIGIT
Uniprot ID	Q495A1
Description	Anti-TIGIT (vibostolimab biosimilar) mAb
Delivery	In Stock
Yefei Storage	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals.
Usage	Research use only

Anti-TIGIT (vibostolimab biosimilar) mAb ELISA

0.2 µg of Human TIGIT, His tagged protein per well




Figure 1. ELISA plate pre-coated by 2 µg/ml (100 µl/well) Human TIGIT, His tagged protein ([getskuurl sku="PME100506"]) can bind Anti-TIGIT Neutralizing antibody in a linear range of 1.6-200.0 ng/ml.






Figure 2. Flow cytometry analysis with 15 µg/mL Anti-TIGIT (vibostolimab) mAb (BME100027) on HEK293 cells transfected with Human TIGIT protein (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

DIMABIO CONFIDENTIAL

