

PRODUCT INFORMATION

Warning: Undefined variable ShasAttributeValueDescription in C:\u00e4wroot\mirror.dimablo.com\wp-content\plugins\woocommerce-print-products-public.php on line 2806
[EN-1029, Hessabody-DhS/DNA, H-DRS-0/JDO.]

Conjugate

Synonyms

Unconjugated TNFRSF10B;TRAILR2;TRAIL-R2;CD262;DR5;KILLER;TRICK2;ZTNFR9;TRICKB

Applications ELISA; Flow Cyt Recommend Dilutions ELISA 1:5000-10000

Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. Formulation & Reconstitution

IgG type Human IgG1 - kappa Reactivity TNFRSF10B Target

Uniprot ID Anti-TNFRSF10B (tigatuzumab biosimilar) mAb Description

Delivery

Background

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.

Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals

Usage

Anti-TNFRSF10B (tigatuzumab biosimilar) mAb ELISA

0.2 µg of Human TNFRSF10B, mFc tagged protein per well





Figure 1. ELISA plate pre-coated by 2 μ g/ml (100 μ l/well) Human TNFRSF10B, mFc tagged protein PME100465 can bind Anti-TNFRSF10B Neutralizing antibody in a linear range of 3.2-80 ng/ml.

Email: info@dimabio.com Website: www.dimabio.com










Figure 2. TNFRSF10B protein is highly expressed on the surface of HEK293 cell membrane. Flow cytometry analysis with Anti-TNFRSF10B (tigatuzumab) on HEK293 cells transfected with human TNFRSF10B(Blue histogram)or HEK293 transfected with irrelevant protein(Red histogram), and Isotype antibody on HEK293 transfected with irrelevant protein(Orange histogram)




Figure 3. Flow cytometry analysis of antigen binding of anti-human TNFRSF10B mAb(BME100032). (A) BME100032 does not bind to TT cells that do not express TNFRSF10B. (B) A clear peak shift of BME100032 was seen compared to the control when incubated with TNFRSF10B-expressing SNU-5 cells, indicating strong binding of BME100032 to TNFRSF10B. Antibodies were incubated at 5 μ g/mL.



Email: info@dimabio.com Website: www.dimabio.com

