

PRODUCT INFORMATION

Warning . Undefined variable \$hasAttributeValueDescription in Calwawroottmirror.dimablo.com/wp-content\plugins/woocommerce-print-products-public.php on line 2806 bM140 Clone ID

TNFSF12 TNFSF12 Host Species

Rabbit Anti-TNFSF12 antibody(DM140); Rabbit mAb Description

Delivery In Stock Uniprot ID 043508 lgG type Rabbit IgG Clonality Monoclonal Reactivity Human ELISA; Flow Cyt Applicati

Recommend Dilutions ELISA 1:5000-10000; Flow Cyt 1:100

Purification Purified from cell culture supernatant by affinity chromatography

Formulation & Reconstitution Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis fo specific instructions of reconstitution.

Storage & Shipping

specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature for the protein and thawing. Lyophilized proteins are shipped at ambient temperature from the protein is a ligand for the five protein encoded by this gene is a cytokine that belongs to the tumor necrosis factor (TNF) ligand mainly. This grotein is a ligand for the exists in both membrane-bound and secreted forms; can induce apoptosis via multiple pattiways of cell death in a cell type-specific manner. This cytokine is also found to promote proliferation and migration of endothelial cells; and thus acts as a regulator of agolgenesis. Alternative splicing results in multiple transcript variants. Some transcripts skip the last exon of this gene and continue into the second exon of the neighboring TNFSF13 gene; such read-through transcripts are contained in Genella Q1977; TNFSF12-TNFSF13.

Usage Research use only

Inconjugated
All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement. DIMA Disclaime




Figure 1. ELISA plate pre-coated by 1 μg/ml (100 μl/well) Human TNFSF12 protein, hFc tagged protein ([getskuurl sku="PME100105"]) can bind Rabbit anti-TNFSF12 monoclonal antibody(clone: DM140) in a linear range of 5-200 ng/ml.

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

Email: info@dimabio.com Website: www.dimabio.com








Figure 2. Flow cytometry analysis with Anti-TNFSF12 (DM140) on HEK293 cells transfected with human TNFSF12 (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

Email: info@dimabio.com Website: www.dimabio.com

