


PRODUCT INFORMATION

Common Name	Warning: Undefined variable \$hasAttributeValueDescription in C:\wwwroot\mirror\dimabio.com\wp-content\plugins\woocommerce-print-products\public\class-woocommerce-print-products-public.php on line 2806 INX 021;INX021;INX-021;SAR-441344;SAR441344;SAR 441344
Conjugate	Biotinylated
Synonyms	IGM;IMD3;TRAP;gp39;CD154;CD40L;HIGM1;T-BAM;TNFSF5;hCD40L
Applications	ELISA
Recommended Dilutions	ELISA 1:5000-10000
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Host Species	N/A
IgG type	IgG1
Reactivity	Human
Target	CD40 Ligand
Uniprot ID	P29965
Description	Biotinylated Anti-CD40 Ligand(frexalimab biosimilar) mAb
Delivery	In Stock
Yefei Storage	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals.
Usage	Research use only

Biotinylated Anti-CD40 Ligand(frexalimab biosimilar) ELISA

0.2 µg of Human CD40 Ligand, hFc tagged protein per well



Biotinylated Anti-CD40 Ligand(frexalimab biosimilar) mAb (ng/mL)

Figure 1. ELISA plate pre-coated by 2 µg/mL (100 µL/well) Human CD40 Ligand Protein, hFc tag (PME100192) can bind Biotinylated Anti-CD40 Ligand(frexalimab biosimilar) mAb (BME100516B) in a linear range of 80-400 ng/mL .

