

PRODUCT INFORMATION

Clone ID	Warning: Undefined variable \$hasAttributeValueDescription in C:\wwwroot\mirror\dimabio.com\wp-content\plugins\woocommerce-print-products\publicclass-woocommerce-print-products-public.php on line 2806 DM191
Target	CD70
Synonyms	CD70;CD27LG;TNFSF7;TNFSF7G;CD27L
Host Species	Rabbit
Description	Biotinylated Anti-CD70 antibody(DM191); Rabbit mAb
Delivery	In Stock
Uniprot ID	P32970
IgG type	Rabbit IgG
Clonality	Monoclonal
Reactivity	Human
Applications	Flow Cyt
Recommended Dilutions	Flow Cyt 15µg/ml
Purification	Purified from cell culture supernatant by affinity chromatography
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Yefel Storage	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	The protein encoded by this gene is a cytokine that belongs to the tumor necrosis factor (TNF) ligand family. This cytokine is a ligand for TNFRSF27;CD27. It is a surface antigen on activated; but not on resting; T and B lymphocytes. It induces proliferation of costimulated T cells; enhances the generation of cytolytic T cells; and contributes to T cell activation. This cytokine is also reported to play a role in regulating B-cell activation; cytotoxic function of natural killer cells; and immunoglobulin synthesis.
Usage	Research use only
Conjugate	Biotinylated
DIMA Disclaimer	All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement.




Figure 1. Flow cytometry analysis with 15µg/ml Biotinylated Anti-CD70 (DM191)mAb on HEK293 cells transfected with Human CD70 protein (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

