PRODUCT INFORMATION

warmon Name
Warning: Undefined variable ShasAttributeValueDescription in C:\wwwroot\mirror.dimabio.com\wp-content\plugins\woocommerce-print-products-public.php on line 2806
18f1, IMC-18f1, L*5012212

 Conjugate
 Biotinylated

 Synonyms
 FLT;FLT-1;VEGFR-1;VEGFR1

 Applications
 ELISA; Flow Cyt

Recommended Dilutions ELISA 1:5000-10000; Flow Cyt 1:100

Formulation & Reconstitution Service from the rile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Host Species Homo sapiens IgG type IgG1
Reactivity Human
Target FLT1

Uniprot ID P17948

Description Biotinylated Anti-FLT1(icrucumab biosimilar) mAb

Delivery 2-3 week

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.

Background Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals

Usage Research use onl

Elisa assay to evaluate Biotinylated Anti-FLT1(icrucumab biosimilar) mAb 0.2µg Human FLT1 Protein, hFc Tag per well




Figure 1. Elisa plates were pre-coated with Human FLT1 Protein, hFc Tag (0.2µg/per well). Serial diluted Biotinylated Anti-FLT1(icrucumab biosimilar) mAb(BME100131B) solutions were added, washed, and incubated with SA-HRP before Elisa reading. From above data, the EC50 for Biotinylated Anti-FLT1(icrucumab biosimilar)monoclonal antibody binding with FLT1 Protein ,hFc Tag is 15.9 ng/ml.

Email: info@dimabio.com

Website: www.dimabio.com

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

