Cat. No. BME100211B



## **PRODUCT INFORMATION**

Warning. Undefined variable \$hasAttributeValueDescription in C:\text{Auwwroot\mirror.dimabio.com\wp-content\plugins\woocommerce-print-products-public.php on line 2806}
Hen egg \text{Vsozyne}
Biotinylated Common Name

Conjugate Synonyms

ELISA, Flow Cyt Recommended Dilutions ELISA 1:5000-10000, Flow Cyt 1:100

Formulation & Reconstitution Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Host Species IgG type IgG2 Reactivity Target HEL Uniprot ID

Applications

Background

Description Biotinylated Anti-HEL Human IgG2-Kappa Isotype control mAb

Delivery

## Control

### Co Biotinylated Anti-HEL Human igls\_Anappa busype town of many 2-3 weeks

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized antibodies are shipped at ambient temperature.

Anti Chicken Hen Egg Lysozyme, specifically recognises Hen Egg Lysozyme (HEL), known also as muramidase or N-acetylmuramide glycanhydrolase, a 14kDa enzymic profein involved in the destruction of bacteria, Lysozyme damages bacterial cell walls by catalog hydrolysis of 1,4e-bet-alinkage, between N-acetylmuramic acid and N-acetyl-D-glucosamine residues in a peptidoglycan and between N-acetyl-D-glucosamine residues in chitodextrins. high concentrations of lysozyme are present in egg white. C-type lysozymes are closely related to a pha-lactalbumin in sequence and structure making them part of the same family.

Research use only

> Email: info@dimabio.com Website: www.dimabio.com

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

