

## PRODUCT INFORMATION

|   |   |
|---|---|
| <b>Clone ID</b>                         | <b>Warning:</b> Undefined variable \$hasAttributeValueDescription in C:\wwwroot\mirror.dimabio.com\wp-content\plugins\woocommerce-print-products\publicclass-woocommerce-print-products-public.php on line 2806<br>1A2  |
| <b>Target</b>                           | IFNG  |
| <b>Synonyms</b>                         | IFN-gamma;Immune interferon   |
| <b>Host Species</b>                     | Rabbit  |
| <b>Description</b>                      | Biotinylated Anti-IFNG antibody(1A2), IgG1 Chimeric mAb   |
| <b>Delivery</b>                         | 2-3 weeks   |
| <b>Uniprot ID</b>                       | P01579  |
| <b>IgG type</b>                         | Rabbit/Human Fc chimeric IgG1   |
| <b>Clonality</b>                        | Monoclonal  |
| <b>Reactivity</b>                       | Human   |
| <b>Applications</b>                     | ELISA   |
| <b>Recommended Dilutions</b>            | ELISA 1/5000-10000  |
| <b>Purification</b>                     | Purified from cell culture supernatant by affinity chromatography   |
| <b>Formulation &amp; Reconstitution</b> | Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.   |
| <b>Yefei_Storage</b>                    | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.   |
| <b>Background</b>                       | IFN $\gamma$ is the major interferon produced by mitogenically or antigenically stimulated lymphocytes. It is structurally different from type I interferon and its major activity is immunoregulation. It has been implicated in the expression of class II histocompatibility antigens in cells that do not normally produce them, leading to autoimmune disease. Interferon gamma is produced mainly by T-cells and natural killer cells activated by antigens, mitogens, or alloantigens. It is produced by lymphocytes expressing the surface antigens CD4 and CD8. IFN $\gamma$ synthesis is induced by IL-2, FGF-basic, and EGF. |
| <b>Usage</b>                            | Research use only   |
| <b>Conjugate</b>                        | Biotinylated  |
| <b>DIMA Disclaimer</b>                  | All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement.  |

DIMABIO CONFIDENTIAL

