

## PRODUCT INFORMATION

|   |  |
|---|--|
| <b>Clone ID</b>                         | <b>Warning:</b> Undefined variable \$hasAttributeValueDescription in C:\wwwroot\mirror.dimabio.com\wp-content\plugins\woocommerce-print-products\publicclass-woocommerce-print-products-public.php on line 2806<br>1A1   |
| <b>Target</b>                           | IL13   |
| <b>Synonyms</b>                         | IL-13:NC30   |
| <b>Host Species</b>                     | Rabbit   |
| <b>Description</b>                      | Biotinylated Anti-IL13 antibody(1A1), IgG1 Chimeric mAb  |
| <b>Delivery</b>                         | 2-3 weeks  |
| <b>Uniprot ID</b>                       | P35225   |
| <b>IgG type</b>                         | Rabbit/Human Fc chimeric IgG1  |
| <b>Clonality</b>                        | Monoclonal   |
| <b>Reactivity</b>                       | Human  |
| <b>Applications</b>                     | ELISA  |
| <b>Recommended Dilutions</b>            | ELISA 1/5000-10000   |
| <b>Purification</b>                     | Purified from cell culture supernatant by affinity chromatography  |
| <b>Formulation &amp; Reconstitution</b> | Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.  |
| <b>Yefei_Storage</b>                    | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.  |
| <b>Background</b>                       | This gene encodes an immunoregulatory cytokine produced primarily by activated Th2 cells. This cytokine is involved in several stages of B-cell maturation and differentiation. It up-regulates CD23 and MHC class II expression, and promotes IgE isotype switching of B cells. This cytokine down-regulates macrophage activity, thereby inhibits the production of pro-inflammatory cytokines and chemokines. This cytokine is found to be critical to the pathogenesis of allergen-induced asthma but operates through mechanisms independent of IgE and eosinophils. This gene, IL3, IL5, IL4, and CSF2 form a cytokine gene cluster on chromosome 5q, with this gene particularly close to IL4. [provided by RefSeq, Jul 2008] |
| <b>Usage</b>                            | Research use only  |
| <b>Conjugate</b>                        | Biotinylated   |
| <b>DIMA Disclaimer</b>                  | All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement.   |

DIMABIO CONFIDENTIAL

