Cat. No. DME100125B



PRODUCT INFORMATION

Warning: Undefined variable ShasAttributeValueDescription in C:\www.root\mirror.dimablo.com\wp-content\plugins\woocommerce-print-products\public\class-woocommerce-print-products-public.php on line 2806 bM125 Clone ID

II 17RΔ

CD217;CDw217;IL-17RA;IL17R;CANDF5;hIL-17R Synonyme

Host Species Rabbit

Biotinylated Anti-IL17RA antibody(DM125); Rabbit mAb Description

Delivery 2-3 weeks Q96F46 Uniprot ID lgG type Rabbit IgG Clonality Monoclonal Reactivity Human Application Flow Cyt Recommended Dilutions Flow Cyt 1:100

Purification Purified from cell culture supernatant by affinity chromatography

Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. Formulation & Reconstitution

Storage & Shipping

Background

specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.

Interleukin 174, (IL-174) is a proinflammatory cyclokine secreted by arctivated 1-1-4/mphocytes. It is a potent inducer of the maturation of CD34-positive hematopoietic precursors into neutrophils. The transmembrane protein encoded by this gene (interleukin 174 Areceptor; IL17RA) is a ubiquitous type I membrane glycoprotein that binds with low affinity to interleukin 174. Interleukin 174 and its receptor play appropriate provided in the provid

Email: info@dimabio.com Website: www.dimabio.com

Usage

Coniugate

Asin seq. All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement. DIMA Disclaimer