Cat. No. DME100126B



PRODUCT INFORMATION

Clone ID DM126 **Target** IL17RA

Synonyms CD217;CDw217;IL-17RA;IL17R;CANDF5;hIL-17R

Host Species Rabbit

Biotinylated Anti-IL17RA antibody(DM126); Rabbit Description

mAb

Delivery 2-3 weeks **Uniprot ID** Q96F46 Rabbit IgG IgG type Clonality Monoclonal Reactivity Human

Applications ELISA; Flow Cyt

Recommended

ELISA 1:5000-10000; Flow Cyt 1:100 **Dilutions**

Purified from cell culture supernatant by affinity **Purification**

chromatography

Lyophilized from sterile PBS, pH 7.4. Normally 5 % Formulation & - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Reconstitution

for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store Storage & Shipping at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

témperature.

Interleukin 17A (IL-17A) is a proinflammatory cytokine secreted by activated T-lymphocytes. It is a potent inducer of the maturation of CD34-

positive hematopoietic precursors into neutrophils. The transmembrane protein encoded

by this gene (interleukin 17A receptor; IL17RA) is a ubiquitous type I membrane glycoprotein that binds with low affinity to interleukin 17A. Interleukin 17A and its receptor play a pathogenic

Background

role in many inflammatory and autoimmune diseases such as rheumatoid arthritis. Like other cytokine receptors; this receptor likely has a multimeric structure. Alternative splicing results in multiple transcript variants encoding different

isoforms.

Usage Research use only

Conjugate Biotinylated

> All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or

> > Email: info@dimabio.com Website: www.dimabio.com

DIMA Disclaimer reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement.

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China)

/+86-400-006-0995(China)

