Cat. No. DMC100276B



## PRODUCT INFORMATION

Warning: Undefined variable shasAttributeValueDescription in Cs\www.root\wirror.dimablo.com\wp-content\plugins\woocommerce-print-products\public.chass-woocommerce-print-products-public.php on line 2806 bMC276 Clone ID

VEGFA MVCD1; VEGF; VPF Synonyme

Host Species Rabbit

Biotinylated Anti-VEGFA antibody(DMC276); IgG1 Chimeric mAb Description

Delivery 2-3 weeks Uniprot ID P15692

IgG type Rabbit/Human Fc chimeric IgG1

Clonality Monoclonal Reactivity Human Application Flow Cyt Recommended Dilutions Flow Cyt 1:100

Purification Purified from cell culture supernatant by affinity chromatography

Formulation & Reconstitution Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Storage & Shipping

Score at 20°C to -80°C for 12 months in Jophilized form. After reconstitution, I not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing), Lophilized proteins are shipped at ambient temperature.

This gene is a member of the PDGFVEG growth factor family, it encodes a heparin-binding protein, which exists as a disuffice-linked homodimer. This process is a member of the PDGFVEG growth factor family, it encodes a heparin-binding protein, which exists as a disuffice-linked homodimer. This process is a member of the PDGFVEG growth factor family, it encodes a heparin-binding protein, which exists as a disuffice-linked homodimer. This process is a member of the PDGFVEG growth factor family. It encodes a heparin-binding protein, which exists as a disuffice-linked homodimer. This process is a support of the PDGFVEG growth factor family. It encodes a heparin-binding protein, which exists as a disuffice-linked homodimer. This process is a support of the PDGFVEG growth factor family. It encodes a heparin-binding protein, which is gene in the process of the PDGFVEG growth factor family. It is expression is correlated with tumor stage and progression. Elevated levels of this protein are found in patients with PDGFVEG growth factor provided growth factor for process of the protein are found in patients with PDGFVEG growth factor growth factor family and progression. Elevated levels of this protein are found in patients with PDGFVEG growth factor growth factor family and progression of on a literative unariative translation initiation of an alternative in-frame translation termination codon via a stop codon readdrough mechanism; and that this isoform is antiangiogenic. Expression of some isoforms derived from the AUG start codon is regulated by a small upstream open reading frame; which is located within an internal ribosome retry site.

Research use only Background

> Email: info@dimabio.com Website: www.dimabio.com

Conjugate

Jon. At Americal Control of the Cont All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement.

