

PRODUCT INFORMATION

CXCR2 **Target**

CD182; IL8R2; IL8RA; IL8RB; CMKAR2; WHIMS2; Synonyms

CDw128b

Recombinant Cynomolgus CXCR2(1-63) protein **Description**

with C-terminal human Fc tag

Delivery In Stock A0A7N9IBZ8 **Uniprot ID Expression Host HEK293**

C-Human Fc tag Tag

Molecular

Purity

Background

CXCR2(Met1-Lys63) hFc(Glu99-Ala330) Characterization

The protein has a predicted molecular mass of **Molecular Weight** 33.4 kDa after removal of the signal peptide.

The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue

staining.

Lyophilized from sterile PBS, pH 7.4. Normally 5 % Formulation & - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Reconstitution

for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store

Storage & Shipping at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

The protein encoded by this gene is a member of the G-protein-coupled receptor family. This protein is a receptor for interleukin 8 (IL8). It binds to IL8 with high affinity, and transduces the signal through a G-protein activated second messenger system. This receptor also binds to chemokine (C-X-C motif) ligand 1 (CXCL1/MGSA), a protein with melanoma growth stimulating activities and has been shown to be a major activity, and has been shown to be a major component required for serum-dependent melanoma cell growth. This receptor mediates

neutrophil migration to sites of inflammation. The angiogenic effects of IL8 in intestinal microvascular endothelial cells are found to be mediated by this receptor. Knockout studies in mice suggested that this receptor controls the positioning of oligodendrocyte precursors in developing spinal cord by arresting their migration. This gene, IL8RA, a gene encoding another high affinity IL8 receptor, as well as IL8RBP, a pseudogene of IL8RB, form a gene cluster in a region mapped to chromosome 2q33q36. Alternatively spliced variants, encoding the same protein, have been identified. [provided by

> Email: info@dimabio.com Website: www.dimabio.com

RefSeq, Nov 2009]

Usage Research use only Conjugate Unconjugated








Figure 1. Cynomolgus CXCR2(1-63) Protein, hFc Tag on SDS-PAGE under reducing condition.

Email: info@dimabio.com Website: www.dimabio.com

