

PRODUCT INFORMATION

| | |
|---|--|
| Target | 5T4 |
| Synonyms | TPBG; M6P1; 5T4AG; WAIF1 |
| Description | Recombinant human 5T4(164-208) Protein with C-terminal human Fc tag |
| Delivery | In Stock |
| Uniprot ID | Q13641 |
| Expression Host | HEK293 |
| Tag | C-Human Fc tag |
| Molecular Characterization | 5T4(Gly164-Ala208) hFc(Glu99-Ala330) |
| Molecular Weight | The protein has a predicted molecular mass of 30.8 kDa after removal of the signal peptide. |
| Purity | The purity of the protein is greater than 90% as determined by SDS-PAGE and Coomassie blue staining. |
| Formulation & Reconstitution | Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. |
| Storage & Shipping | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | This gene encodes a leucine-rich transmembrane glycoprotein that may be involved in cell adhesion. The encoded protein is an oncofetal antigen that is specific to trophoblast cells. In adults this protein is highly expressed in many tumor cells and is associated with poor clinical outcome in numerous cancers. Alternate splicing in the 5' UTR results in multiple transcript variants that encode the same protein. [provided by RefSeq, Oct 2009] |
| Usage | Research use only |
| Conjugate | Unconjugated |






Figure 1. Human 5T4(164-208) Protein, hFc Tag on SDS-PAGE under reducing condition.

DIMABIO CONFIDENTIAL

