

PRODUCT INFORMATION

Tag	C-Flag Tag
Target	ADGRE2
Synonyms	CD97; CD312; EMR2; VBU
Description	Human ADGRE2 full length protein-synthetic nanodisc
Delivery	In Stock
Uniprot ID	Q9UHX3
Expression Host	HEK293
Protein Families	Druggable Genome, GPCR, Transmembrane
Protein Pathways	N/A
Molecular Weight	The human full length ADGRE2 protein has a MW of 90.5 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	This gene encodes a member of the class B seven-span transmembrane (TM7) subfamily of G-protein coupled receptors. These proteins are characterized by an extended extracellular region with a variable number of N-terminal epidermal growth factor-like domains coupled to a TM7 domain via a mucin-like spacer domain. The encoded protein is expressed mainly in myeloid cells where it promotes cell-cell adhesion through interaction with chondroitin sulfate chains. This gene is situated in a cluster of related genes on chromosome 19. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.
Usage	Research use only
Conjugate	Unconjugated



ELISA assay to evaluate ADGRE2-Nanodisc 0.2 μ g Human ADGRE2-Nanodisc per well




Figure1. Elisa plates were pre-coated with Flag Tag ADGRE2-Nanodisc (0.2 μ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with ADGRE2-Nanodisc is 3.100ng/ml.



Figure2. Human ADGRE2-Nanodisc, Flag Tag on SDS-PAGE

