

PRODUCT INFORMATION

Warning . Undefined variable \$hasAttributeValueDescription in Calwawroottmirror.dimablo.com/wp-content\plugins/woocommerce-print-products/public/class-woocommerce-print-products-public.php on line 2806 CFlagStrps To. Tag

ADORA2R

Synonyme

Human ADORA2B-Strep full length protein-synthetic nanodisc

Description Delivery 6~8weeks Uniprot ID P29275 Expression Host HEK293

Protein Families Druggable Genome, GPCR, Transmembrane

Protein Pathways Calcium signaling pathway, Neuroactive ligand-receptor interaction, Vascular smooth muscle contraction

Molecular Weight The human full length ADORA2B-Strep protein has a MW of 36.3 kDa

Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. Formulation & Reconstitution

Storage & Shipping

DIMARIO CONTINUE DE LA CONTINUE DE L ryop-mization, recease see Letrincate or Analysis for Specific Instructions of reconstitution. Store at -20°C to -80°C for 12 months in Jophilized form. After reconstitution, in fon intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. This gene encodes an adenosine receptor that is a member of the G protein-coupled receptor superfamily. This integral membrane protein stimulates adenylate cyclose activity in the presence of adenosine. This protein also interacts with netrin-1, which is involved in axon elongation. The gene is located near the Smith-Magenis syndrome region on chromosome 1.7. (provided by RefSeq. jul 2008)

> Email: info@dimabio.com Website: www.dimabio.com

Usage Research use only Conjugate

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

