

PRODUCT INFORMATION

AGTR2 **Target**

Synonyms AT2; ATGR2; MRX88

Recombinant human AGTR2 Protein with C-Description

terminal human Fc tag

Delivery In Stock **Uniprot ID** P50052 **Expression Host HEK293**

Tag C-Human Fc tag

Molecular

Purity

Background

AGTR2(Met1-Asp45) hFc(Glu99-Ala330) Characterization

The protein has a predicted molecular mass of **Molecular Weight** 30.9 kDa after removal of the signal peptide.

The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue

staining.

Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before

Formulation & lyophilization. Please see Certificate of Analysis Reconstitution for specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Storage & Shipping

Lyophilized proteins are shipped at ambient

temperature.

The protein encoded by this gene belongs to the G-protein coupled receptor 1 family, and functions as a receptor for angiotensin II. It is an intergral membrane protein that is highly expressed in fetus and in neonates, but scantily

in adult tissues, except brain, adrenal medulla, and atretic ovary. This receptor has been shown to mediate programmed cell death and this apoptotic function may play an important role in developmental biology and pathophysiology. Mutations in this gene are been associated with X-linked cognitive disability. Severe Acute

Respiratory Syndrome Coronavirus (SARS-CoV) and SARS-CoV-2 infection results in downregulation of angiotensin converting enzyme-2

(ACE2) receptors, the effects of which, triggers serious inflammatory lesions in the tissues involved, primarily in the lungs. The inflammatory reaction appears to be mediated by angiotensin II


derivatives, including the angiotensin AT2 receptor which has been found to be upregulated

> Email: info@dimabio.com Website: www.dimabio.com

in bronchoalveolar lavage samples from Coronavirus disease 2019 (COVID19) patients. [provided by RefSeq, Jul 2020]

Usage Research use only

Conjugate Unconjugated










Figure 1. Human AGTR2 Protein, hFc Tag on SDS-PAGE under reducing condition.

