

PRODUCT INFORMATION

Target	Beta crystallin S
Synonyms	CRYG8; CTRCT20
Description	Recombinant protein of human crystallin, gamma S (CRYGS)
Delivery	2-3 weeks
Uniprot ID	P22914
Expression Host	HEK293T
Tag	C-Myc/DDK
Molecular Characterization	N/A
Molecular Weight	20.8 kDa
Purity	> 80% as determined by SDS-PAGE and Coomassie blue staining
Formulation & Reconstitution	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10% glycerol
Storage & Shipping	Store at -80°C.
Background	<p>Crystallins are separated into two classes: taxon-specific, or enzyme, and ubiquitous. The latter class constitutes the major proteins of vertebrate eye lens and maintains the transparency and refractive index of the lens. Since lens central fiber cells lose their nuclei during development, these crystallins are made and then retained throughout life, making them extremely stable proteins. Mammalian lens crystallins are divided into alpha, beta, and gamma families; beta and gamma crystallins are also considered as a superfamily. Alpha and beta families are further divided into acidic and basic groups. Seven protein regions exist in crystallins: four homologous motifs, a connecting peptide, and N- and C-terminal extensions. Gamma-crystallins are a homogeneous group of highly symmetrical, monomeric proteins typically lacking connecting peptides and terminal extensions. They are differentially regulated after early development. This gene encodes a protein initially considered to be a beta-crystallin but the encoded protein is monomeric and has greater sequence similarity to other gamma-crystallins. This gene encodes the most significant gamma-crystallin in adult eye lens tissue. Whether due to aging or mutations in specific genes, gamma-crystallins have been involved in cataract formation. [provided by RefSeq, Jul 2008]</p>
Usage	Research use only
Conjugate	Unconjugated

