

**PRODUCT INFORMATION**

|   |   |
|---|---|
| <b>Target</b>                           | CA12  |
| <b>Synonyms</b>                         | CAXII; CA-XII; T18816; HsT18816   |
| <b>Description</b>                      | Recombinant human CA12 Protein with C-terminal 10×His tag   |
| <b>Delivery</b>                         | In Stock  |
| <b>Uniprot ID</b>                       | O43570  |
| <b>Expression Host</b>                  | HEK293  |
| <b>Tag</b>                              | C-10×His tag  |
| <b>Molecular Characterization</b>       | CA12(Ala25-Ser301) 10×His tag   |
| <b>Molecular Weight</b>                 | The protein has a predicted molecular mass of 32.5 kDa after removal of the signal peptide. The apparent molecular mass of CA12-His is approximately 25-55 kDa due to glycosylation.  |
| <b>Purity</b>                           | The purity of the protein is greater than 85% as determined by SDS-PAGE and Coomassie blue staining.  |
| <b>Formulation &amp; Reconstitution</b> | Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.  |
| <b>Storage &amp; Shipping</b>           | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.   |
| <b>Background</b>                       | Carbonic anhydrases (CAs) are a large family of zinc metalloenzymes that catalyze the reversible hydration of carbon dioxide. They participate in a variety of biological processes, including respiration, calcification, acid-base balance, bone resorption, and the formation of aqueous humor, cerebrospinal fluid, saliva, and gastric acid. This gene product is a type I membrane protein that is highly expressed in normal tissues, such as kidney, colon and pancreas, and has been found to be overexpressed in 10% of clear cell renal carcinomas. Three transcript variants encoding different isoforms have been identified for this gene. [provided by RefSeq, Jun 2014] |
| <b>Usage</b>                            | Research use only   |
| <b>Conjugate</b>                        | Unconjugated  |






Figure 1. Human CA12 Protein, His Tag on SDS-PAGE under reducing condition.

DIMABIO CONFIDENTIAL

