

## **PRODUCT INFORMATION**

Warning: Undefined variable \$hasAttributeValueDescription in C:\www.root\mirror.dimablo.com\wp-content\plugins\woocommerce-print-products-public.php on line 2806 C-Flag Tag Tag

CANN6; CB-R; CNR1; CB1A; CB1K5; CB1R; CNR Human CB1 full length protein-synthetic nanodisc

Delivery Uniprot ID P21554 HEK293 **Expression Host** Protein Families GPCR

Protein Pathways Neuroactive ligand-receptor interaction

Molecular Weigh The human full length CB1 Protein has a MW of 52.7 kDa

Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. Formulation & Reconstitution

Storage & Shipping

lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. Store at -20° C to -80°C for 12° months in lyophilized form. After reconstitution, in for intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.

The cannabinoids, principally delate-3-tetrahydrocannabinol and synthetic analogs, are psychoactive ingredients of marijuana. The cannabinoid receptors are members of the guanine-nucleotide-binding protein (G-protein) coupled receptor family, which inhibit adenylate cyclase activity in a dose-dependent, stereoselective and pertusiss toxin-sensitive manner. The two receptors have been found to be invoided in the cannabinoid-induced CNS effects (including alterations in mood and cognition) experienced by users of marijuana. Multiple transcript variants encoding two different protein isoforms have been described for this gene. Background

Usage Research use only Unconjugated

## ELISA assay to evaluate CB1-Nanodisc 0.2µg Human CB1-Nanodisc per well




Figure 1. Elisa plates were pre-coated with Flag Tag CB1-Nanodisc ( $0.2\mu g/per$  well). Serial diluted anti-CB1 monoclonal antibody (DME100144) solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-CB1 monoclonal antibody binding with CB1-Nanodisc is 41.62ng/ml.

Email: info@dimabio.com Website: www.dimabio.com








Figure 2. Human CB1-Nanodisc, Flag Tag on SDS-PAGE



Email: info@dimabio.com Website: www.dimabio.com