

## **PRODUCT INFORMATION**

CCL5 **Target** 

C-C Motif Chemokine 5;EoCP;Eosinophil

Chemotactic Cytokine;SIS-Delta;Small-Inducible Cytokine A5;T Cell-Specific Protein P228;TCP228;T-Cell-Specific Protein RANTES;CCL5;D17S136E;SCYA5 Synonyms

Recombinant Human C-C Motif Chemokine 5 is produced by our E.coli expression system and the target gene encoding Ser24-Ser91 is expressed. **Description** 

**Delivery** In Stock **Uniprot ID** P13501 **Expression Host** E.coli

Tag

Molecular Characterization

Not available

Molecular Weight 7.8 KDa

Greater than 95% as determined by reducing **Purity** 

SDS-PAGE.

Lyophilized from a 0.2  $\mu m$  filtered solution of 20mM PB, 150mM NaCl, pH 7.4. Formulation &

Reconstitution

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store Storage & Shipping

at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

témperature.

Human Chemokine (C-C Motif) Ligand 5 (CCL5) plays an active role in recruiting leukocytes into inflammatory sites. CCL5 is secreted by many cell types at inflammatory sites and it exerts a wide range of activities through the receptors CCR1, CCR3, CCR4, and CCR5. N-Terminal truncated CCL5/RANTES, Met-RANTES, and amino-oxygentane (AOP)-RANTES exhibit antagonist or partial against functions on their recentors.

**Background** 

partial agonist functions on their receptors. CCL5/RANTES attracts different subtypes of leukocytes into inflamed tissue and intervenes in

> Email: info@dimabio.com Website: www.dimabio.com

a wide range of allergic and autoimmune

diseases.

Research use only **Usage** 

Conjugate Unconjugated








Figure 1. Greater than 95% as determined by reducing SDS-PAGE.



Email: info@dimabio.com Website: www.dimabio.com