

PRODUCT INFORMATION

CD117 **Target**

Synonyms PBT; SCFR; C-Kit; KIT; MASTC

Recombinant human CD117 Protein with C-Description

terminal mouse Fc tag

Delivery In Stock **Uniprot ID** P10721 **Expression Host HEK293**

Tag C-Mouse Fc tag

Molecular

Purity

Background

CD117(Glu925-Val976) mFc(Pro99-Lys330) Characterization

The protein has a predicted molecular mass of **Molecular Weight** 31.9 kDa after removal of the signal peptide.

The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue

staining.

Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before

Formulation & lyophilization. Please see Certificate of Analysis Reconstitution

for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Storage & Shipping

Lyophilized proteins are shipped at ambient

temperature.

This gene encodes a receptor tyrosine kinase. This gene was initially identified as a homolog of the feline sarcoma viral oncogene v-kit and is often referred to as proto-oncogene c-Kit. The

canonical form of this glycosylated

transmembrane protein has an N-terminal extracellular region with five immunoglobulin-like domains, a transmembrane region, and an intracellular tyrosine kinase domain at the Cterminus. Upon activation by its cytokine ligand, stem cell factor (SCF), this protein phosphorylates multiple intracellular proteins that play a role in in the proliferation, differentiation, migration and apoptosis of many cell types and thereby plays an important role in homotopicies from cell

important role in hematópoiesis, stem céll maintenance, gametogenesis, melanogenesis, and in mast cell development, migration and function. This protein can be a membrane-bound or soluble protein. Mutations in this gene are

associated with gastrointestinal stromal tumors, mast cell disease, acute myelogenous leukemia, and piebaldism. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, May 2020]


> Email: info@dimabio.com Website: www.dimabio.com

Usage Research use only Conjugate Unconjugated

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China)

/+86-400-006-0995(China)





JE. Figure 1. Human CD117 Protein, mFc Tag on SDS-PAGE under reducing condition.



