

PRODUCT INFORMATION

ributeValueDescription in C:\wwwroot\mirror.dimabio.com\wp-content\plugins\woocommerce-print-rce-print-products-public.php on line 2806 Tag

Human CHRM2 full length protein-synthetic nanodisc

Delivery Uniprot ID P08172 HEK293 **Expression Host**

Protein Families Druggable Genome, GPCR, Transmembrane

Protein Pathways Calcium signaling pathway, Neuroactive ligand-receptor interaction, Regulation of actin cytoskeleton

Molecular Weigh The human full length CHRM2 protein has a MW of 51.7 kDa

Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. Formulation & Reconstitution

lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Store at 2-0°C to 80°C for 12 cmonths in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.

The muscarinic cholinergic receptors belong to a larger family of 6 protein-coupled receptors. The functional diversity of these receptors is defined by the binding of acetylcholine to these receptors and includes cellular responses such as adenylate cyclase inhibition, phosphoinositide degeneration, and potassium channel mediation. Muscarnic receptors influence many effects of acetylcholine in the central and peripheral nervous system. The muscarnic cholinergic receptor 2 is involved in mediation of bradycardia and a decrease in cardiac contractility. Multiple alternatively spliced transcript variants have been described for this gene, [provided by Neffsec, jul 2008]

Research use only Unconjugated

Background

ELISA assay to evaluate CHRM2-Nanodisc 0.2µg Human CHRM2-Nanodisc per well




Figure 1. Elisa plates were pre-coated with Flag Tag CHRM2-Nanodisc (0.2µg/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with CHRM2-Nanodisc is 13.52ng/ml.

Email: info@dimabio.com Website: www.dimabio.com










Figure 2. Human CHRM2-Nanodisc, Flag Tag on SDS-PAGE



Email: info@dimabio.com Website: www.dimabio.com