

PRODUCT INFORMATION

Target	Cyclin D2
Synonyms	KIAK0002; MPPH3
Description	Recombinant protein of human cyclin D2 (CCND2)
Delivery	2-3 weeks
Uniprot ID	P30279
Expression Host	HEK293T
Tag	C-Myc/DDK
Molecular Characterization	N/A
Molecular Weight	32.9 kDa
Purity	> 80% as determined by SDS-PAGE and Coomassie blue staining
Formulation & Reconstitution	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10% glycerol
Storage & Shipping	Store at -80°C.
Background	<p>The protein encoded by this gene belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance through the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. This cyclin forms a complex with CDK4 or CDK6 and functions as a regulatory subunit of the complex, whose activity is required for cell cycle G1/S transition. This protein has been shown to interact with and be involved in the phosphorylation of tumor suppressor protein Rb. Knockout studies of the homologous gene in mouse suggest the essential roles of this gene in ovarian granulosa and germ cell proliferation. High level expression of this gene was observed in ovarian and testicular tumors. Mutations in this gene are associated with megalencephaly-polymicrogyria-polydactyly-hydrocephalus syndrome 3 (MPPH3). [provided by RefSeq, Sep 2014]</p>
Usage	Research use only
Conjugate	Unconjugated

