

PRODUCT INFORMATION

Target FCGR3A

FCGR3A;CD16A;FCG3;FCGR3;IGFR3 **Synonyms**

Recombinant human FCGR3A protein(F176V) with Description

C-terminal 6×His tag

Delivery In Stock **Uniprot ID** P08637 **Expression Host HEK293** Tag C-6×His Tag

Molecular

FCGR3A(Gly17-Gly206)(F176V) 6×His tag Characterization

The protein has a predicted molecular mass of **Molecular Weight** 22.3 kDa after removal of the signal peptide.

The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue

Purity

staining.

Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before Formulation & Reconstitution

lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Storage & Shipping

Lyophilized proteins are shipped at ambient

temperature.

This gene encodes a receptor for the Fc portion of immunoglobulin G, and it is involved in the removal of antigen-antibody complexes from the circulation, as well as other other antibody-dependent responses. This gene (FCGR3A) is highly similar to another nearby gene (FCGR3B) located on chromosome 1. The receptor encoded by this gene is expressed on natural killer (NK) cells as an integral membrane glycoprotein anchored through a transmembrane peptide,

Background whereas FCGR3B is expressed on

polymorphonuclear neutrophils (PMN) where the

receptor is anchored through a phosphatidylinositol (PI) linkage. Mutations in this


gene have been linked to susceptibility to recurrent viral infections, susceptibility to systemic lupus erythematosus, and alloimmune neonatal neutropénia. Alternatively spliced transcript variants encoding different isoforms

have been found for this gene.


Usage Research use only

Conjugate Unconjugated










15 Figure 1. Human FCGR3A Protein (F176V), His Tag on SDS-PAGE under reducing condition.

Human FCGR3A(F176V), His Tagged protein ELISA

0.2 μg of Human FCGR3A(F176V), His tagged protein per well



, ...,

Figure 2. ELISA plate pre-coated by 2 μ g/mL (100 μ L/well) Human FCGR3A Protein (F176V), His Tag (PME100069) can bind Anti-FCGR3A(AFM13 without CD30 biosimilar) mAb (BME100267) in a linear range of 0.64–80 ng/mL.








Human FCGR3A(F176V), His Tagged protein ELISA

0.2 μg of Human FCGR3A(F176V), His tagged protein per well



Anti-FCGR3A(AFM13_VH_2 without CD30 biosimilar) mAb (ng/mL)

Figure 3. ELISA plate pre-coated by 2 μ g/mL (100 μ L/well) Human FCGR3A Protein (F176V), His Tag (PME100069) can bind Anti-FCGR3A(AFM13_VH_2 without CD30 biosimilar) mAb (BME100270) in a linear range of 3.20–400 ng/mL.

Email: info@dimabio.com Website: www.dimabio.com

