

PRODUCT INFORMATION

Warning: Undefined variable ShasAttributeValueDescription in C:\www.root\mirror.dimabio.com\wp-content\plugins\woocommerce-print-products-public.php on line 2806
C-FlagdStrep Tag Tag

FML2_HUMAN, FMLP-R-II, FMLPY, FPRH1, FPRH2, FPRL2, RMLP-R-I Synonyme Human FPR3-Strep full length protein-synthetic nanodisc

Description Delivery 6~8weeks Uniprot ID P25089

Expression Host HEK293 Protein Families GPCR,Transmembrane,Druggable Genome GPCRDB Class A Rhodopsin-like,Peptide GPCRs, Protein Pathways

Molecular Weight

The human full length FPR3-Strep protein has a MW of 40 kDa Lyophilized from ananoids: solubilization buffer (20 mM Tris-HCI. 150 mM NaCI, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. Formulation & Reconstitution

Storage & Shipping

OINIABIO CONTINUINALIA DINIABIO lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. Store at -20°C to -80°C for 12 cmonths in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.

Low affinity receptor for N-formyl-methionly peoplitics, which are powerful neutrophils chemotactic factors. Binding of FMLP to the receptor causes activation of neutrophils. This response is mediated via a G-protein that activates a phosphatidylinositol-calcium second messenger system.[UniProtKl5/wiss-Prot Function]

Research use only

> Email: info@dimabio.com Website: www.dimabio.com

Usage Conjugate Unconjugated

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

