

## **PRODUCT INFORMATION**

Warning . Undefined variable \$hasAttributeValueDescription in Calwawroottmirror.dimablo.com/wp-content\plugins/woocommerce-print-products/public/class-woocommerce-print-products-public.php on line 2806 CFlag Tag Tag

GLP-1; GLP-1-R; GLP-1R

Human GLP1R full length protein-synthetic nanodisc

Delivery Uniprot ID P43220 HEK293 **Expression Host** 

Protein Families Druggable Genome, ES Cell Differentiation/IPS, GPCR, Transmembra

Protein Pathways Neuroactive ligand-receptor interaction

Molecular Weigh The human full length GLP1R protein has a MW of 53.0 kDa

Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. Formulation & Reconstitution

Storage & Shipping

lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.

A 7-transmembrane protein that functions as a receptor for glucagon-like peptide 1 (GLP-1) hormone, which stimulates glucose-induced insulin secretion. This receptor, which functions at the cell surface, becomes internalized in response to GLP-1 and GLP-1 analogs, and it plays an important role in the signaling cascades leading to insulin secretion. It also displays neuroprotective effects in animal models, Polymorphisms in this gene are associated with diabetes. The protein is an important drug target for the treatment of type 2 diabetes and stroke. Alternative splicing of this gene results in multiple transcript variants. Background

Research use only

Usage Unconjugated

## ELISA assay to evaluate GLP1R-Nanodisc 0.2µg Human GLP1R-Nanodisc per well




Figure 1. Elisa plates were pre-coated with Flag Tag GLP1R-Nanodisc ( $0.2\mu g/per$  well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with GLP1R-Nanodisc is 3.549ng/ml.

Email: info@dimabio.com Website: www.dimabio.com










Figure 2. Human GLP1R-Nanodisc, Flag Tag on SDS-PAGE



Email: info@dimabio.com Website: www.dimabio.com