

PRODUCT INFORMATION

| | |
|---|---|
| Tag | C-Flag&Strep Tag |
| Target | GPR18 |
| Synonyms | N-arachidonyl glycine receptor; NAGly receptor; |
| Description | G-protein coupled receptor 18 GPCRW GPR18 Human GPR18-Strep full length protein-synthetic nanodisc |
| Delivery | 6~8weeks |
| Uniprot ID | Q14330 |
| Expression Host | HEK293 |
| Protein Families | GPCR,Transmembrane,Druggable Genome, |
| Protein Pathways | GPCRDB Other, |
| Molecular Weight | The human full length GPR18-Strep protein has a MW of 38.1 kDa Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments. |
| Formulation & Reconstitution | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Storage & Shipping | |
| Background | Receptor for endocannabinoid N-arachidonyl glycine (NAGly) (PubMed:16844083, PubMed:24762058, PubMed:27572937). However, conflicting results about the role of NAGly as an agonist are reported (PubMed:27018161). Can also be activated by plant-derived and synthetic cannabinoid agonists (PubMed:24762058). The activity of this receptor is mediated by G proteins which inhibit adenylyl cyclase (PubMed:16844083). May contribute to regulation of the immune system. Is required for normal homeostasis of CD8 subsets of intraepithelial lymphocytes (IELs) (CD8alphaalpha and CD8alphabeta IELs)in small intestine by supporting preferential migration of CD8alphaalpha T-cells to intraepithelial compartment over lamina propria compartment, and by mediating their reconstitution into small intestine after bone marrow transplant (By similarity). Plays a role in hypotensive responses, mediating reduction in intraocular and blood pressure (By similarity). Mediates NAGly-induced process of reorganization of actin filaments and induction of acrosomal exocytosis (PubMed:27572937).[UniProtKB/Swiss-Prot Function] |
| Usage | Research use only |
| Conjugate | Unconjugated |

