

PRODUCT INFORMATION

Target	HRG
Synonyms	HPRG; HRGP; THPH11
Description	Recombinant protein of human histidine-rich glycoprotein (HRG)
Delivery	2-3 weeks
Uniprot ID	P04196
Expression Host	HEK293T
Tag	C-Myc/DDK
Molecular Characterization	N/A
Molecular Weight	57.6 kDa
Purity	> 80% as determined by SDS-PAGE and Coomassie blue staining
Formulation & Reconstitution	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10% glycerol
Storage & Shipping	Store at -80°C.
Background	This histidine-rich glycoprotein contains two cystatin-like domains and is located in plasma and platelets. The physiological function has not been determined but it is known that the protein binds heme, dyes and divalent metal ions. The encoded protein also has a peptide that displays antimicrobial activity against <i>C. albicans</i> , <i>E. coli</i> , <i>S. aureus</i> , <i>P. aeruginosa</i> , and <i>E. faecalis</i> . It can inhibit rosette formation and interacts with heparin, thrombospondin and plasminogen. Two of the protein's effects, the inhibition of fibrinolysis and the reduction of inhibition of coagulation, indicate a potential prothrombotic effect. Mutations in this gene lead to thrombophilia due to abnormal histidine-rich glycoprotein levels. [provided by RefSeq, Nov 2014]
Usage	Research use only
Conjugate	Unconjugated

