

PRODUCT INFORMATION

LIF **Target**

Leukemia inhibitory factor; D factor; **Synonyms**

MLPLI;Emfilermin

Recombinant human LIF protein with C-terminal **Description**

human Fc tag

Delivery In Stock P15018 **Uniprot ID Expression Host** HFK293

Tag C-Human Fc Tag

Molecular

Storage & Shipping

Background

LIF(Ser23-Phe202) hFc(Glu99-Ala330) Characterization

The protein has a predicted molecular mass of 46 kDa after removal of the signal peptide. The apparent molecular mass of LIF-hFc is **Molecular Weight**

approximately 55-70 kDa due to glycosylation. The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue

Purity

staining.

Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before Formulation & lyophilization. Please see Certificate of Analysis Reconstitution

for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient

temperature.

The protein encoded by this gene is a pleiotropic cytokine with roles in several different systems. It is involved in the induction of hematopoietic differentiation in normal and myeloid leukemia cells, induction of neuronal cell differentiation, regulator of mesenchymal to epithelial conversion

during kidney development, and may also have a role in immune tolerance at the maternal-fetal interface. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq, Mar 2012]

Usage Research use only

Conjugate Unconjugated

> Email: info@dimabio.com Website: www.dimabio.com








Figure 1. Human LIF Protein, hFc Tag on SDS-PAGE under reducing condition.

Human LIF, hFc Tagged protein ELISA

 $0.2~\mu g$ of Human LIF, hFc tagged protein per well




Figure 2. ELISA plate pre-coated by 2 μ g/mL (100 μ L/well) Human LIF Protein, hFc Tag (PME100744) can bind Anti-LIF(falbikitug biosimilar) mAb (BME100506) in a linear range of 2.29–61.73 ng/mL, mouse anti-human Fab-specific antibody was used as detection antibody.



