

PRODUCT INFORMATION

Warning: Undefined variable \$hasAttributeValueDescription in C:\www.root\mirror.dimablo.com\wp-content\plugins\woocommerce-print-products\public\class-woocommerce-print-products-public.php on line 2806 C-flag Tag Tag

EDG7, Edg-7, GPCR, HOFNH30, LP-A3, LPA3, RP4-678I3 Synonyme Human LPAR3 full length protein-synthetic nanodisc

Description Delivery 6~8weeks Uniprot ID O9UBY5 HEK293 Expression Host

Protein Families GPCR, Transmembrane, Druggable Genome Protein Pathways G-Protein Coupled Receptors Signaling Pathway, Molecular Weight The human full length LPAR3 protein has a MW of 40.1kDa

Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCI, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. Formulation & Reconstitution

Storage & Shipping

DIMARIO CONTINUE DE LA CONTINUE DE L yopminication: recease see Lettricate or Analysis for Specific Instructions of reconstitution. Store at -20°C to -80°C for 12 months in Jopolinized form. After reconstitution, in for intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. This gene encodes a member of the G protein-coupled receptor family, as well as the EDG family of proteins. This protein functions as a cellular receptor for lysophosphatidic acid and mediates lysophosphatidic acid-evoked calcium mobilization. This receptor couples predominantly to Gig/11) alpha proteins, provided by Refese, Jul 2008]

> Email: info@dimabio.com Website: www.dimabio.com

Usage Research use only Conjugate

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

