

PRODUCT INFORMATION

Warning: Undefined variable \$hasAttributeValueDescription in C:\www.root\mirror.dimabio.com\wp-content\plugins\woocommerce-print-products-public.php on line 2806 C-HagdStrep Tag Tag

BLT2, BLTR2, JULF2, KPG_004, LTB4-R 2, LTB4-R2, NOP9 Synonyme Human LT4R2-Strep full length protein-synthetic nanodisc

Description Delivery 6~8weeks Uniprot ID O9NPC1 HEK293 Expression Host

Protein Families

GPCR,Transmembrane,Druggable Genome,
GPCRDB Other,Asthma,Autoimmune & Inflammatory Response, Protein Pathways Molecular Weight

The human full length LT4R2-Strep protein has a MW of 37.9 kDa
Lyophilized from nanodics solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before
lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. Formulation & Reconstitution

Storage & Shipping

DIMARIO CONTINUE DE LA CONTINUE DE L ryopimization. recease see Letinicate or Analysis for Specific Instructions of reconstitution. Store at -20°C to 80°C for 12 months in lyophilized form. After reconstitution, in for intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.

Low-affinity receptor for leukorienes including leukorteine BA. Mediates chemotaxis of granulocytes and macrophages. The response is mediated via G-proteins that activate a phosphatidylinositol-calcium second messenger system. The rank order of affinities for the feukotrienes is LTB4 > 12-epi-LTB4 > LTB3 > LTB3. | LTB3. |

> Email: info@dimabio.com Website: www.dimabio.com

Usage Research use only Conjugate



