

## **PRODUCT INFORMATION**

N-MBP Tag, C-Flag Tag Tag

**Target** CLDN6

**Synonyms** Claudin 6, Claudin-6, Skullin, Claudin 6

Human MBP-CLDN6 full length protein-synthetic **Description** 

nanodisc In Stock

**Delivery Uniprot ID** P56747 **Expression Host HEK293** 

Formulation &

Storage & Shipping

**Background** 

Usage

**Protein Families** Transmembrane

Cell adhesion molecules (CAMs), Leukocyte **Protein Pathways** transendothelial migration, Tight junction

The human full length MBP-CLDN6 Protein has a **Molecular Weight** 

MW of 63.3 kDa

Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis

Reconstitution for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store

at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

Tight junctions represent one mode of cell-to-cell adhesion in epithelial or endothelial cell sheets, forming continuous seals around cells and serving as a physical barrier to prevent solutes and water from passing freely through the paracellular space. These junctions are comprised of sets of continuous networking strands in the outwardly facing cytoplasmic leaflet, with complementary grooves in the inwardly facing extracytoplasmic

leaflet. This gene encodes a component of tight junction strands, which is a member of the claudin family. The protein is an integral membrane protein and is one of the entry cofactors for hepatitis C virus. The gene methylation may be involved in esophageal tumorigenesis. This gene is adjacent to another family member CLDN9 on chromosome 16.

> Email: info@dimabio.com Website: www.dimabio.com

Research use only

Conjugate Unconjugated

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China)

/+86-400-006-0995(China)





## ELISA assay to evaluate MBP-CLDN6-Nanodisc 0.2µg Human MBP-CLDN6-Nanodisc per well




Figure 1. Elisa plates were pre-coated with N-MBP Tag, C-Flag Tag MBP-CLDN6-Nanodisc ( $0.2\mu g/per$  well). Serial diluted anti-CLDN6 monoclonal antibody (BME100082) solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-CLDN6 monoclonal antibody binding with MBP-CLDN6-Nanodisc is 723.6ng/ml.

Email: info@dimabio.com Website: www.dimabio.com




Figure 2. Human MBP-CLDN6-Nanodisc with N-MBP Tag, C-Flag Tag on SDS-PAGE

