

PRODUCT INFORMATION

Tag	Warning: Undefined variable \$hasAttributeValueDescription in C:\wwwroot\mirror.dimabio.com\wp-content\plugins\woocommerce-print-products\public\class-woocommerce-print-products-public.php on line 2806 C-Flag&Strep Tag
Target	NPFF2
Synonyms	GPR74, HLWAR77, NPFF2, NPGPR
Description	Human NPFF2-Strep full length protein-synthetic nanodisc
Delivery	6~8weeks
Uniprot ID	Q9Y5X5
Expression Host	HEK293
Protein Families	Transmembrane,Druggable Genome,
Protein Pathways	GPCRDB Class A Rhodopsin-like,Neurotrophin & Receptors,
Molecular Weight	The human full length NPFF2-Strep protein has a MW of 60.3 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	This gene encodes a member of a subfamily of G-protein-coupled neuropeptide receptors. This protein is activated by the neuropeptides A-18-amide (NPAF) and F-8-amide (NPFF) and may function in pain modulation and regulation of the opioid system. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jan 2009]
Usage	Research use only
Conjugate	Unconjugated

