

PRODUCT INFORMATION

| | |
|---|--|
| Tag | C-Flag Tag |
| Target | NPSR1 |
| Synonyms | ASRT2; GPR154; GPRA; NPSR; PGR14; VRR1 |
| Description | Human NPSR1 full length protein-synthetic nanodisc |
| Delivery | In Stock |
| Uniprot ID | Q6W5P4 |
| Expression Host | HEK293 |
| Protein Families | Druggable Genome, Transmembrane |
| Protein Pathways | N/A |
| Molecular Weight | The human full length NPSR1 protein has a MW of 42.7 kDa |
| Formulation & Reconstitution | Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments. |
| Storage & Shipping | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | A member of the vasopressin/oxytocin subfamily of G protein-coupled receptors. The encoded membrane protein acts as a receptor for neuropeptide 5 and affects a variety of cellular processes through its signaling. Increased expression of this gene in ciliated cells of the respiratory epithelium and in bronchial smooth muscle cells is associated with asthma. Polymorphisms in this gene have also been associated with asthma susceptibility, panic disorders, inflammatory bowel disease, and rheumatoid arthritis. Alternative splicing results in multiple transcript variants. |
| Usage | Research use only |
| Conjugate | Unconjugated |



ELISA assay to evaluate NPSR1-Nanodisc 0.2 μ g Human NPSR1-Nanodisc per well




Figure1. Elisa plates were pre-coated with Flag Tag NPSR1-Nanodisc (0.2 μ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with NPSR1-Nanodisc is 6.386ng/ml.

kDa M R




Figure2. Human NPSR1-Nanodisc, Flag Tag on SDS-PAGE

