

PRODUCT INFORMATION

Tag C-Flag&Strep Tag

Target OPRM

Synonyms LMOR; M-OR-1; MOP; MOR; MOR1; OPRM1

DescriptionHuman OPRM-Strep full length protein-synthetic

nanodisc 6~8weeks

Delivery 6~8weeks
Uniprot ID P35372
Expression Host HEK293

Storage & Shipping

Background

Protein Families Druggable Genome, GPCR, Transmembrane

Protein Pathways Neuroactive ligand-receptor interaction

Molecular Weight

The human full length OPRM-Strep protein has a

MW of 44.8 kDa Lyophilized from nanodisc solubilization buffer (20

mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5%
– 8% trehalose is added as protectants before
lyophilization. Please see Certificate of Analysis
for specific instructions. Do not use solvents with

for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient

temperature.

One of at least three opioid receptors in humans; the mu opioid receptor (MOR). The MOR is the principal target of endogenous opioid peptides and opioid analgesic agents such as betaendorphin and enkephalins. The MOR also has an important role in dependence to other drugs of abuse, such as nicotine, cocaine, and alcohol via

endorphin and enkephalins. The MOR also has ar important role in dependence to other drugs of abuse, such as nicotine, cocaine, and alcohol via its modulation of the dopamine system. The NM_001008503.2:c.118A>G allele has been associated with opioid and alcohol addiction and variations in pain sensitivity but evidence for it

transcript variants encoding different isoforms have been found for this gene. Though the canonical MOR belongs to the superfamily of 7-transmembrane-spanning G-protein-coupled receptors some isoforms of this gene have only 6

having a causal role is conflicting. Multiple

transmembrane domains.

Usage Research use only
Conjugate Unconjugated

Email: info@dimabio.com Website: www.dimabio.com



