

PRODUCT INFORMATION

\$hasAttributeValueDescription in C:\wwwroot\mirror.dimabio.com\wp-content\plugins\woocommerce-print-ommerce-print-products-public.php on line 2806 Tag

ORXR2; OX2R; OXR2

Human HCRTR2-Strep full length protein-synthetic nanodisc

Delivery Uniprot ID 043614 HEK293 **Expression Host**

Protein Families Druggable Genome, GPCR, Transmembrane Protein Pathways Neuroactive ligand-receptor interaction

Molecular Weigh The human full length HCRTR2-Strep protein has a MW of 50.7 kDa

Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCI, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. Formulation & Reconstitution Storage & Shipping

syopimization. Presse see Celtificate or Airospas for specific instructions or reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.

A G-protein coupled receptor involved in the regulation of feeding behavior. The encoded protein binds the hypothalamic neuropeptides orexin A and orexin B. A related gene (HCRTR1) encodes a G-protein coupled receptor that selectively binds orexin A. Background Research use only

Conjugate Unconjugated

ELISA assay to evaluate HCRTR2-Strep-Nanodisc 0.2µg Human HCRTR2-Strep-Nanodisc per well




Figure 1. Elisa plates were pre-coated with C-Flag&Strep Tag HCRTR2-Strep-Nanodisc (0.2µg/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with HCRTR2-Strep-nanodisc is 2.093ng/ml.

Email: info@dimabio.com Website: www.dimabio.com










Figure 2. Human HCRTR2-Strep-Nanodisc, C-Flag&Strep Tag on SDS-PAGE



