Cat. No. FLP120429

Delivery

Formulation &

Background



PRODUCT INFORMATION

C-Flag&Strep Tag Tag

OXYR Target

OT-R; OTR **Synonyms**

Human OXYR-Strep full length protein-synthetic **Description**

nanodisc 6~8weeks P30559

Uniprot ID Expression Host HEK293

Protein Families Druggable Genome, GPCR, Transmembrane

Calcium signaling pathway, Neuroactive ligand-**Protein Pathways**

receptor interaction

The human full length OXYR-Strep protein has a **Molecular Weight**

MW of 42.8 kDa

Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis

Reconstitution for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store Storage & Shipping at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

The protein belongs to the G-protein coupled receptor family and acts as a receptor for oxytocin. Its activity is mediated by G proteins

which activate a phosphatidylinositol-calcium second messenger system. The oxytocin-oxytocin receptor system plays an important role in the

Email: info@dimabio.com Website: www.dimabio.com

uterus during parturition.

Usage Research use only Conjugate Unconjugated

