

PRODUCT INFORMATION

iable \$hasAttributeValueDescription in Ct\wwwroot\mirror.dimabio.com\wp-content\plugins\woocommerce-print-woocommerce-print-products-public.php on line 2806 Tag

AGR16; DFNB68; EDG-5; EDG5; Gpcr13; H218; LPB2; S1P2 Human S1PR2-Strep full length protein-synthetic nanodisc

Delivery Uniprot ID 095136 HEK293 **Expression Host**

Protein Families Druggable Genome, GPCR, Transmembrane Neuroactive ligand-receptor interaction Protein Pathways

Molecular Weigh The human full length S1PR2-Strep protein has a MW of 38.9 kDa

Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCI, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. Formulation & Reconstitution Storage & Shipping

Store at .20°C to .80°C for 12 months in lyophilized form. After reconstitution, if not interned for use within a month, aliquot and store at -80°C (Avoid Receptor for the lyosphin

Research use only Usage Conjugate Unconjugated

ELISA assay to evaluate S1PR2-Strep-Nanodisc 0.2µg Human S1PR2-Strep-Nanodisc per well




Figure 1. Elisa plates were pre-coated with C-Flag&Strep Tag S1PR2-Strep-Nanodisc (0.2 μ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with S1PR2-Strep-nanodisc is 1.269ng/ml.

Email: info@dimabio.com Website: www.dimabio.com

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)






Figure 2. Human S1PR2-Strep-Nanodisc, C-Flag&Strep Tag on SDS-PAGE

