

PRODUCT INFORMATION

Tag	C-Flag Tag
Target	SCARB1
Synonyms	CD36L1;CLA-1;CLA1;HDLQTL6;SR-BI;SRB1
Description	Human SCARB1 full length protein-synthetic nanodisc
Delivery	In Stock
Uniprot ID	Q8WTV0
Expression Host	HEK293
Protein Families	Druggable Genome, Transmembrane
Protein Pathways	N/A
Molecular Weight	The human full length SCARB1 protein has a MW of 60.9 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	The protein is a plasma membrane receptor for high density lipoprotein cholesterol (HDL). The encoded protein mediates cholesterol transfer to and from HDL. In addition, this protein is a receptor for hepatitis C virus glycoprotein E2. Several transcript variants encoding different isoforms have been found for this gene.
Usage	Research use only
Conjugate	Unconjugated



ELISA assay to evaluate SCARB1-Nanodisc 0.2 μ g Human SCARB1-Nanodisc per well




Figure1. Elisa plates were pre-coated with Flag Tag SCARB1-Nanodisc (0.2 μ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with SCARB1-Nanodisc is 8.388ng/ml.



Figure2. Human SCARB1-Nanodisc, Flag Tag on SDS-PAGE

