

PRODUCT INFORMATION

Tag	Warning: Undefined variable \$hasAttributeValueDescription in C:\wwwroot\mirror.dimabio.com\wp-content\plugins\woocommerce-print-products\public\class-woocommerce-print-products-public.php on line 2806 C-Flag Tag
Target	SCNBA
Synonyms	FEPS3, HSNAN7, NAV1.9, NaN, PMS, SCN12A, SNS-2
Description	Human SCNBA full length protein-synthetic nanodisc
Delivery	6-8weeks
Uniprot ID	Q9UI33
Expression Host	HEK293
Protein Families	Ion Channels: Sodium
Protein Pathways	N/A
Molecular Weight	The human full length SCNBA protein has a MW of 204.9kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	Voltage-gated sodium channels are transmembrane glycoprotein complexes composed of a large alpha subunit with 24 transmembrane domains and one or more regulatory beta subunits. They are responsible for the generation and propagation of action potentials in neurons and muscle. This gene encodes one member of the sodium channel alpha subunit gene family, and is highly expressed in nociceptive neurons of dorsal root ganglia and trigeminal ganglia. It mediates brain-derived neurotrophic factor-evoked membrane depolarization and is a major effector of peripheral inflammatory pain hypersensitivity. Mutations in this gene have been associated with hereditary sensory and autonomic neuropathy type VII and familial episodic pain syndrome-3. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Mar 2017]
Usage	Research use only
Conjugate	Unconjugated

