

PRODUCT INFORMATION

| | |
|---|--|
| Tag | C-Flag Tag |
| Target | SCTR |
| Synonyms | SR |
| Description | Human SCTR full length protein-synthetic nanodisc |
| Delivery | 6~8weeks |
| Uniprot ID | P47872 |
| Expression Host | HEK293 |
| Protein Families | Transmembrane,Druggable Genome, |
| Protein Pathways | GPCRDB Class B Secretin-like,Cancer, |
| Molecular Weight | The human full length SCTR protein has a MW of 50.2kDa |
| Formulation & Reconstitution | Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments. |
| Storage & Shipping | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | The protein encoded by this gene is a G protein-coupled receptor and belongs to the glucagon-VIP-secretin receptor family. It binds secretin which is the most potent regulator of pancreatic bicarbonate, electrolyte and volume secretion. Secretin and its receptor are suggested to be involved in pancreatic cancer and autism. [provided by RefSeq, Jul 2008] |
| Usage | Research use only |
| Conjugate | Unconjugated |

