

PRODUCT INFORMATION

Tag	Warning: Undefined variable \$hasAttributeValueDescription in C:\wwwroot\mirror.dimabio.com\wp-content\plugins\woocommerce-print-products\public\class-woocommerce-print-products-public.php on line 2806 C-Flag&Strep Tag
Target	STEAP2
Synonyms	IPCA1; PCANAP1; PUMPCn; STAMP1; STMP
Description	Human STEAP2-Strep full length protein-synthetic nanodisc
Delivery	In Stock
Uniprot ID	Q8NFT2
Expression Host	HEK293
Protein Families	Transmembrane
Protein Pathways	N/A
Molecular Weight	The human full length STEAP2-Strep protein has a MW of 56.1 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. A member of the STEAP family and encodes a multi-pass membrane protein that localizes to the Golgi complex, the plasma membrane, and the vesicular tubular structures in the cytosol. A highly similar protein in mouse has both ferriredutase and cupric reductase activity, and stimulates the cellular uptake of both iron and copper in vitro. Increased transcriptional expression of the human gene is associated with prostate cancer progression. Alternate transcriptional splice variants, encoding different isoforms, have been characterized.
Background	
Usage	Research use only
Conjugate	Unconjugated

ELISA assay to evaluate STEAP2-Strep-Nanodisc
0.2µg Human STEAP2-Strep-Nanodisc per well




Figure 1. Elisa plates were pre-coated with C-Flag&Strep Tag STEAP2-Strep-Nanodisc (0.2µg/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with STEAP2-Strep-nanodisc is 1.357ng/ml.






Figure 2. Human STEAP2-Strep-Nanodisc, C-Flag&Strep Tag on SDS-PAGE

