

## **PRODUCT INFORMATION**

Warning: Undefined variable ShasAttributeValueDescription in C:\www.root\mirror.dimablo.com\wp-content\plugins\woocommerce-print-products\public\class-woocommerce-print-products-public.php on line 2806 C-Flag Tag Tag

ERIS; hMITA; hSTING; MITA; MPYS; NET23; SAVI; STING; STING-beta; TMEM173

Human STING1 full length protein-synthetic nanodisc

Delivery Uniprot ID 086WV6 HEK293 **Expression Host** Protein Families

Protein Pathways Cytosolic DNA-sensing pathway, RIG-I-like receptor signaling pathway

Molecular Weigh The human full length STING1 protein has a MW of 42.2 kDa

Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCI, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. Formulation & Reconstitution

Storage & Shipping

yopnization, rease see Letrincate or Analysis for specim instructions or reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.

It is a pattern recognition receptor that detects, cytosolic nucleic acids and transmits signals that activate type I interferon responses. The encoded protein has also been shown to play a role in apoptotic signaling by associating with type II major histocompatibility complex. Mutations in this gene are the cause of infantile-lowest STIMS—associated vasculopathy. Alternate splicing results in multiple transcript variants.

Conjugate Unconjugated

## ELISA assay to evaluate STING1-Nanodisc 0.2µg Human STING1-Nanodisc per well




Figure 1. Elisa plates were pre-coated with Flag Tag STING1-Nanodisc ( $0.2\mu g/per$  well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with STING1-Nanodisc is 5.896 ng/ml.

Email: info@dimabio.com Website: www.dimabio.com










Figure 2. Human STING1-Nanodisc, Flag Tag on SDS-PAGE



Email: info@dimabio.com Website: www.dimabio.com