

## **PRODUCT INFORMATION**

Warning: Undefined variable ShasAttributeValueDescription in C\u00e4wroot\u00fcmirror.dimabio.com\u00e4wp-content\plugins\u00e4woocommerce-print-products\u00e4public\class-woocommerce-print-products-public.php on line 2806 C-flag Tag Tag

Synonyme

Human SUCR1 full length protein-synthetic nanodisc 6~8weeks

Description Delivery Uniprot ID O9BXA5 HEK293 Expression Host

Protein Families GPCR,Transmembrane,Druggable Genome

GPCRDB Class A Rhodopsin-like, Protein Pathways

Molecular Weight The human full length SUCR1 protein has a MW of 38.7kDa

DIMARIO CONTINUE DE LA CONTINUE DE L Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCI, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. Formulation & Reconstitution

> Email: info@dimabio.com Website: www.dimabio.com

Storage & Shipping

yopminization: receive see Lettinicate or inanitysis for specific instructions or reconstitution. Store at -20°C to -80°C for 12 months in Joynhilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.

This gene encodes a G-protein-coupled receptor for succinate, an intermediate molecule of the citric acid cycle, It is involved in the promotion of hematopoietic progenitor cell development, and it has a potential role in renovascular hypertension which has known correlations to renal failure, diabetes and atherosclerosis. [provided by ReFSeq, Oct 2009]

Usage Research use only Conjugate



