

**PRODUCT INFORMATION**

<b>Target</b>	TIGIT
<b>Synonyms</b>	TIGIT;VSIG9;VSTM3
<b>Description</b>	Recombinant human TIGIT Protein with C-terminal human Fc tag
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	Q495A1
<b>Expression Host</b>	HEK293
<b>Tag</b>	C-Human Fc Tag
<b>Molecular Characterization</b>	TIGIT(Met22-Pro141) hFc(Glu99-Ala330)
<b>Molecular Weight</b>	The protein has a predicted molecular mass of 39.13 kDa after removal of the signal peptide. The apparent molecular mass of TIGIT-hFc is approximately 40-53 kDa due to glycosylation.
<b>Purity</b>	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Yefei_Storage</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	This gene encodes a member of the PVR (poliovirus receptor) family of immunoglobulin proteins. The product of this gene is expressed on several classes of T cells including follicular B helper T cells (TFH). The protein has been shown to bind PVR with high affinity, this binding is thought to assist interactions between TFH and dendritic cells to regulate T cell dependent B cell responses.
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated



Figure 1. Human TIGIT Protein, hFc Tag on SDS-PAGE under reducing condition.



## Human TIGIT, hFc Tagged protein ELISA

0.2  $\mu$ g of TIGIT, hFc Tagged protein per well




Figure 2. ELISA plate pre-coated by 2  $\mu$ g/ml (100  $\mu$ l/well) Human CD155, mFc-His tagged protein ([getskuurl sku="PME100027"]) can bind Human TIGIT, hFc tagged protein (PME100107) in a linear range of 1.95-125 ng/ml.




Figure 3. Flow cytometry analysis with 1  $\mu$ g/mL Human TIGIT Protein, hFc tag (PME100107) on HEK293 cells transfected with human CD112 (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).



### Human TIGIT, hFc Tagged protein ELISA

0.2  $\mu\text{g}$  of Human TIGIT, hFc tagged protein per well




Figure 4. ELISA plate pre-coated by 2  $\mu\text{g}/\text{mL}$  (100  $\mu\text{L}/\text{well}$ ) Human TIGIT Protein, hFc Tag (PME100107) can bind Anti-TIGIT (etigilimab biosimilar) mAb (BME100024) in a linear range of 0.32–8 ng/mL.

