

PRODUCT INFORMATION

\$hasAttributeValueDescription in C:\wwwroot\mirror.dimabio.com\wp-content\plugins\woocommerce-printcommerce-print-products-public.php on line 2806 Tag

ANKTM1; FEPS; FEPS1

Human TRPA1 full length protein-synthetic nanodisc

Delivery Uniprot ID 075762 HEK293 **Expression Host**

Protein Families Druggable Genome, Ion Channels: Transient receptor potential, Transmembrane

Protein Pathways

Background

Molecular Weigh The human full length TRPA1 protein has a MW of 127.5 kDa

Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. Formulation & Reconstitution

Storage & Shipping

lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Store at -20°C to 80°C for 12 cmonths in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.

The structure of the protein is highly related to both the protein ankyrin and transmembrane proteins. This protein is activated by a large variety of structurally unrelated electrophilic and non-electrophilic chemical compounds. Electrophilic ligands activate TRPA1 by interacting with critical N-terminal Cys residues in a covalent manner, whereas mechanisms of non-electrophilic ligands are not well determined. May be a component for the mechanosensitive transduction channel of hair cells in inner ear, thereby participating in the perception of sounds. Probably operated by a phosphatidy industois lescond messenger system.

Research use only

Unconjugated

ELISA assay to evaluate TRPA1-Nanodisc 0.2µg Human TRPA1-Nanodisc per well




Figure 1. Elisa plates were pre-coated with Flag Tag TRPA1-Nanodisc ($0.2\mu g/per$ well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with TRPA1-Nanodisc is 7.433ng/ml.



Email: info@dimabio.com Website: www.dimabio.com








Figure 2. Human TRPA1-Nanodisc, Flag Tag on SDS-PAGE



Email: info@dimabio.com Website: www.dimabio.com